

PRECONCENTRATION AND DETERMINATION OF NICKEL IONS USING SILICA WITH FUNCTIONAL GROUPS OF SULFONIC DERIVATIVES OF NITROSO NAPHTHOLS

A.K. TROFIMCHUK, S.L. DIDUKH, A.N. MUKHINA, V.N. LOSEV

References:

1. Zambelli B., Ciurli S. Nickel and human health. *Met. Ions. Life Sci.* 2013. 13. 321-357.
2. Duda-Chodak A., Blaszczyk U. The impact of nickel on human health. *J. Elementol.* 2008. 13(4). 685-696.
3. Denkhaus E., Salnikow K. Nickel essentiality, toxicity, and carcinogenicity. *Crit. Rev. Oncol. Hematol.* 2002. 42(1). 35-56.
4. Mirabi A., Jamali M.R., Meheraeeen P., Berijani K. Determination of trace amounts of nickel in food and environmental water samples by flame atomic absorption Spectrometry after dispersive liquid-liquid microextraction. *Asian J. Chem.* 2012. 24(8). 3425-3429.
5. Sánchez-Rojas F., Bosch-Ojeda C. Preconcentration of nickel in waters by vortex assisted dispersive liquid-liquid microextraction. *Sample Prep.* 2017, 3(1). 11-13.
6. Mirzaei M., Behzadi M., Mahmoud Abadi N., Beizaei A. Simultaneous separation/preconcentration of ultra trace heavy metals in industrial wastewaters by dispersive liquid-liquid microextraction based on solidification of floating organic drop prior to determination by graphite furnace atomic absorption spectrometry. *J. Hazard. Mater.* 2011. 186. 1739-1743.
7. Safavi A., Abdollahi H., Hormozi Nezhad M.R., Kamali R., Cloud point extraction preconcentration and simultaneous spectrophotometric determination of nickel and cobalt in water samples. *Spectrochim. Acta.* 2004. 60A. 2897-2901.
8. Kaya G., Akdeniz I., Yaman M. Solid phase extraction and preconcentration of nickel in water samples with activated carbon-PAR and flame atomic absorption spectrometry. *Atomic Spectrosc.* 2008. 29(4). 150-155.
9. Pourreza N., Keykhaii M., Behdarvand N. Solid phase extraction of nickel as methylthymol blue complex on naphthalene adsorbent and flame atomic absorption spectrometric determination. *Canad. J. Analyt. Sci. Spectrosc.* 2006, 51(1), 17-23.
10. Khadem M., Shahtaheri S.J., Golbabaei F., Rahimi-Froushani A. Comparative study of nickel recoveries from urine, nail, and hair samples using XAD-7 and Chromosorb 105 resins. *Inter. J. Occupational Hygiene.* 2014. 6(3). 101-107.
11. Jiang N., Chang X., Zheng H., He Q., Hu Zh. Selective solid-phase extraction of nickel(II) using a surface-imprinted silica gel sorbent. *Analyt. Chim. Acta.* 2006. 577. 225-231.
12. Taher M.A. Preconcentration of trace nickel with the ion pair of disodium 1-nitroso-2-naphthol-3,6-disulfonate and tetradecyldimethylbenzylammonium chloride on microcrystalline naphthalene or by the column method and determination by third derivative spectrophotometry. *Talanta.* 1999. 50(3). 559-567.
13. Amin A.S., AL-Attas A.S. Study of the solid phase extraction and spectrophotometric determination of nickel using 5-(40-chlorophenylazo)-6-hydroxypyrimidine-2,4-dione in environmental samples. *J. Saudi Chem. Soc.* 2012. 16. 451-459.
14. Hu Q., Yang G., Huang Zh., Yin J. Study on solid phase extraction and spectrophotometric determination of nickel in waters and biological samples. *Bull. Korean Chem. Soc.* 2004, 25(4), 545-548.
15. Gazda D.B., Fritz J.S., Porter M.D. Determination of nickel(II) as the nickel dimethylglyoxime complex using colorimetric solid phase extraction. *Analyt. Chim. Acta.* 2004, 508, 53-59.
16. Taher M.A., Dehzoei A.M., Puri B., Puri S. Derivative UV-visible spectrophotometric determination of nickel in alloys and biological samples after preconcentration with the ion pair of 2-nitroso-1-naphthol-4-sulfonic acid and tetradecyldimethylbenzylammonium chloride onto microcrystalline naphthalene or by a column method. *Analyt. Chim. Acta.* 1998. 367(1-3). 55-61.
17. Toel K., Motomizu Sh., Yokosu H. Exrtraction-spectrophotometric determination of nickel with 4-chloro-2-nitrosonaphtol and crystal violet. *Analyt. Chim. Acta.* 1979. 110. 324-334.
18. Ivanov V.M., Samarina T.O., Figurivskaya V.N. Investigation of complex formation of nickel(II) with 1-nitroso-2-naphthol-3,6-disulfonic acid by optical and chromatcity methods. *Vestnik Moskovskogo Universiteta* [Bulletin of the Moscow University]. 2011, 52(4), 285-290 (in Russian).
19. Didukh S.L. Adsorption-photometric determination of iron using silica with nitroso-R salt and nitroso-N salt functional groups. *J. Analyt. Chem.* 2017, 72(1), 49-55. doi: 10.1134/S1061934817010051.
20. Losev, V.N., Didukh, S.L., Mukhina, A.N., Trofimchuk, A.K. Using silica modified by polyhexamethylene guanidine and nitroso-R salt for the preconcentration and determination of cobalt. *J. Analyt. Chem.* 2015, 70(6), 677-684. doi: 10.1134/S106193481506006
21. Akolo S.A., Kovo A.S. Comparative Study of Adsorption of Copper Ion onto Locally Developed and Commercial Chitosan. *J. Encapsul. Ads. Sci.* 2015. 5(1). 21-37.

22. Reshetnyak E., Ivchenko N., Nikitina N. Photometric determination of aqueous cobalt(II), nickel(II), copper(II) and iron(III) with 1-nitroso-2-naphthol-3,6-disulfonic acid disodium salt in gelatin films. *Centr. Europ. J. Chem.* 2012. 10(5). 1617-1623.
23. Azarova Zh.M., Morosanova E.I., Zolotov Yu.A. Xerogels modified with 1-(2-pyridylazo)-2-naphthol and dimethylglyoxime: Indicator tubes for determining nickel. *J. Analyt. Chem.*, 2000, 55(7), 641-644.
24. Dmitrienko S.G., Kosyreva O.A., Runov V.K., Zolotov Yu.A. Utilization of Polyurethane Foams in Sorption-Photometric Analysis. *Mend. Commun.* 1991. 1(2). 75-77.