

Spectrophotometric Analysis of Quaternary Drug Mixtures using Artificial Neural network model

Azhar S. Hamody[†], Faeza H. Zankanah[‡], Saad A. Ali[†], Nahla Alassaf^{†*}, and Sarmad B. Dikran^{***}

[†] Department of Chemistry, College of Education for Pure Science/ Ibn Al-Haitham, Adhamiya, University of Baghdad, Baghdad-Iraq; *e-mail: nahla.a.a@ihcoedu.uobaghdad.edu.iq

[‡] Department of Prosthodontics Technology, College of Health & Medical Technology, Uruk University, Baghdad-Iraq;

^{***} This paper is dedicated to the memory of the late Sarmad B. Dikran. Deceased July 6, 2020;

Received: July 16, 2022; Accepted: September 06, 2022

DOI: 10.17721/moca.2022.118-124

Copyright © 2022 Azhar S. Hamody et al. Open access article distributed under the Creative Commons Attribution License CC BY 4.0.

A Novel artificial neural network (ANN) model was constructed for calibration of a multivariate model for simultaneously quantitative analysis of the quaternary mixture composed of carbamazepine, carvedilol, diazepam, and furosemide. An eighty-four mixing formula were prepared and analyzed spectrophotometrically. Each analyte was formulated in six samples at different concentrations thus twenty-four samples for the four analytes were tested. A neural network of 10 hidden neurons was capable to fit data 100%. The suggested model can be applied for the quantitative chemical analysis for the proposed quaternary mixture.

Keywords: artificial neural network, simultaneous spectrophotometric analysis, quaternary mixture, carbamazepine, carvedilol, diazepam, furosemide

Carvedilol (CAV), Diazepam (DIA) and Furosemide (FUR) are different classes of medications that are used in combination for management of high blood pressure and heart diseases. CAV [(2RS)-1-(9HCarbazol-4-yloxy)-3-[[2-(2-methoxyphenoxy) ethyl] amino] propan-2-ol], nonselective beta-blocker/alpha-blocker drug, commonly applied to control hypertension and treatment of congestive heart failure and cardiac arrhythmia [1, 2]. DIA [7-chloro-1- methyl-5-phenyl-1,3-dihydro-2H-1,4- benzodiazepin-2-one] is

a drug used to manage a variety of medical situations like anxiety and status epilepticus [1,3]. FUR [4-Chloro-2-[(furan-2-ylmethyl) amino] - 5-sulfamoyl benzoic acid], a loop diuretics, commonly used in the management of hypertension, fluid buildup as well as edema [1,4]. CAB [5H-dibenzo [b, f] azepine-5-carboxamide], antiseizure drug, mainly used to treat epilepsy and neuropathic pain [1,4]. The chemical structures [1] of CAB, CAV, DIA and FUR are shown in Figure (1).




Figure 1: Chemical structures of studied drugs.

Literature survey reveals several conventional analytical techniques including HPLC [5-8], FT-IR [9], flow injection analysis [10,11], GC-Mass [12,13], voltammetry [14,15], spectrofluorimetry [16], and spectrophotometry [17-19] for individual or simultaneous quantification of the cited drugs in bulk, pharmaceutical preparations, and body fluids.

The current study aims to evaluate the quaternary synthetic mixture that composed of CAB, CAV, DIA, and FUR in their mixtures via artificial neural networks (ANN). ANN model has the pattern recognition capabilities of the neural networks of the brain and have been used in many pharmaceutical researches [20]. The applications of ANN to quantitative chemical analysis using multivariate instrumental signals with the mathematical aspects involved in the training of neural networks has been reviewed [21]. The principle and theory of the suggested model were discussed in the literature [22-28].

Experimental part

Instruments and software

Shimadzu 1800 UV - Visible Spectrophotometer equipped with 10 mm quartz cells with bundled software version (UV Prob 2.34) was used for recording absorption spectra. The range of the scans was from

(200 – 350) nm at data interval of 0.5 nm.

A Simplex Lattice Mixture Design (create by JMP® 11.0.0 SAS Institute Inc.) was used to prepare a set of calibration mixtures for the analysis of the mentioned drugs simultaneously.

Chemicals and reagents

The used raw materials/powders of CAB, CAV, DIA, and FUR were of pharmaceutical grade with purity of 99.99%, which were kindly obtained as a gift from the State Company for Drug Industries and Medical Appliances Samara-Iraq (SDI), while methanol was of analytical grade reagent.

Standard and working solutions

Standard stock solutions (1000 µg/mL) of CAB, CAV, DIA, and FUR were separately prepared by dissolving 0.050 gm of the cited drugs in 50 mL methanol. Serial dilution was done for the preparation of working solutions of each drug.

Data analysis

All obtained data were analysed using MATLAB (version 9.1.0 R2021a, Math Work, Inc).

Results and Discussion

The normalized absorption spectra of CAB, CAR, DIA, and FUR are shown in Figures (2-5) at concentration range (1-20) µg/mL for each component.




Figure 2: CAB absorbance vs. wavelength




Figure 3: CAV absorbance vs. wavelength




Figure 4: DIA absorbance vs. wavelength




Figure 5: FUR absorbance vs. wavelength

The UV spectra of the examined drugs showed a noteworthy overlap (Figure 6) that hampers the application of the traditional spectrophotometric methods for settling their spectra. In this manner, artificial neural network (ANN) model was used for the concurrent investigation of this complex quaternary mixture.

Eighty-four samples were prepared in order to optimize, calibrate, and validate the suggested model. The utilized absorbance information of the scanned samples was in the range (200 – 350) nm, (Figure 7). The wavelengths lower than 205.5 nm, were omitted because of the effect of high noise, for this reason,

the data that was obtained out of the specified range would have resulted in a substantial noise level within the calibration matrix, thus influencing the precision.

Artificial Neural Network (ANN) model:

Calibration procedure

Neural network could be considered as an excellent multivariate method in modeling the nonlinear behavior such as multicomponent system in chemical analysis. The aim of multivariate method is to construct a calibration between the concentration of the samples and the absorbance. The first step in model is the design of the calibration network for the quaternary complex mixture.




Figure 6: UV absorption spectra of CAB, CAR, DIA, and FUR.




Figure 7: Absorbance spectra of 84 samples of CAB, CAR, DIA, and FUR mixtures in methanol.

The network was configured for 10 neurons in the hidden layer, two input vectors (absorbances and wavelengths) and 4 output vectors (concentration of CAB, CAV, DIA and FUR), the neural network was configured for 290 inputs (absorbance at 205.5 nm – 350 nm). The constructed neural network is illustrated in Figure (8), which demonstrates the training algorithms, the training progress performance.




Figure 8: The neural network used for training the experimental results.

While, Figure (9), shows the best validation found at epoch 12.




Figure 9: The mean square error versus training epoch.

Results shown in Figure (10) illustrated excellent fitting via statistically R values for training, validation, and testing. One of the important points during calculation mode is to reject and retrain any unsatisfied results.




Figure 10: R values for training, validation, testing of the neural network.

Statistically, the MSE and R values for trained, validated, and tested data are shown in Table (1), the total number of trained samples is 108 to get 4 outputs.

For conformation of the ANN model, the data were tested in the model, it showed approximately 100% fitting after retraining. The architecture of the ANN model was plotted as revealed in the Figures (11-14).

Table 1. MSE and R- values for the training data, validity and test data.

Data	Samples	MSE	R
Train	70%	1.0244E-5	0.9999
Valid	15%	0.6040	0.9882
Test	15%	0.8812	0.9886




Figure 11: ANN test for Carbamazepine.




Figure 12: ANN test for Carvedilol.




Figure 13: ANN test for Diazepam.




Figure 14: ANN test for Furosemide.

Conclusion

The results of the current study apply evidence to the scientific data regarding the efficiency of the ANN model in simultaneous estimation of quaternary system with high or distinctive overlapped spectra. Adding to that, this method could be suggested for using in quality control without intrusion, with considering it as a smart method for chemometric analysis.

Reference

1. British Pharmacopoeia (2013), volume I & II, p716, 267, 494, 287.
2. P. Stafylas, P. Sarafidis (2008) Carvedilol in Hypertension Treatment, *Vasc Health Risk Manag*, 4(1): 23-30.
3. N. Calcaterra, J. Barrow (2014) Classics in Chemical Neuroscience: Diazepam (Valium), *ACS chemical neuroscience*, 5(4): 253-260.
4. 19th edition WHO Models List of Essential Medicines (2015) Explanatory notes.
5. S. Dikran, A. Mohammed, N. Alassaf (2016) Simple RP-HPLC Method for Estimation of Furosemide, Carbamazepine, Diazepam and Carvedilol in Bulk and Pharmaceutical Dosage Forms, *Chemistry and Materials Research*, 8(3): 54-60.
6. V. Ram, P. Dave, H. Joshi (2012) Development and Validation of a Stability- Indicating HPLC Assay Method for Simultaneous Determination of Spironolactone and Furosemide in Tablet Formulation, *Journal of Chromatographic Science JCS*, 50: 721-726.
7. B. Böger et al.; (2018) Determination of carbamazepine and diazepam by SPE-HPLC-DAD in Belém River water, Curitiba-PR/Brazil, *Rev. Ambient. Água*, 13(2), e2196.
8. A. Elezovic et al.; (2015) Development and Comparison of Two HPLC Methods, Chiral and Achiral, for Determination of Carvedilol Content in Tablets, *Pharmacia*, 18(1): 30-35.
9. P. Garbacz and M. Wesolowski (2018) DSC, FTIR and Raman Spectroscopy Coupled with Multivariate Analysis in a Study of Co-Crystals of Pharmaceutical Interest, *Molecules*, 23, 2136.
10. P. Biparva, S. Abedirad, S. Kazemi (2014) ZnO Nanoparticles as an Oxidase Mimic-Mediated Flow-Injection Chemiluminescence System for Sensitive Determination of Carvedilol, *Talanta*, 130: 116-121.
11. S. Han, S. Jia, L. Guo (2013) Flow-Injection Chemiluminescence Determination of Diazepam by Oxidation with N-Bromosuccinimide, *Luminescence*, 28(6): 888-893.
12. E. Hoehn (2014) Detection of the Pharmaceuticals Carbamazepine and Diphenhydramine in Tissue Extracts Using Gas Chromatography-Mass Spectrometry (GC-MS), *Environmental Studies Undergraduate Student Theses, Bachelor of Science, University of Nebraska-Lincoln*.
13. O. Zaporozhets, I. Tsyrlneva, M. Ischenko (2012) Determination of 8 Diuretics and Probenecid in Human Urine by Gas Chromatography-Mass Spectrometry: Confirmation Procedure, *American Journal of Analytical Chemistry*, 3: 320- 327.
14. M. Pan et al.; (2014) Determination of Carbamazepine: A Comparison of the Differential Pulse Voltammetry (DPV) Method and the Immunoassay Method in a Clinical Trial, *Journal of Analytical Chemistry*, 69(1): 57- 61.
15. K. Honeychurch et al.; (2013) The Redox Behaviors of Diazepam (Valium®) using a Disposable Screen- Printed Sensor and its Determination in Drinks using a Novel Adsorptive Stripping Voltammetric Assay, *Talanta*, 116: 300- 307.
16. L. Abdel Fattah, T. Mohamed, E. Taha (2010) Spectrofluorimetric Determination of Carvedilol in Dosage Form and Spiked Human Plasma through Derivatization with 1-Dimethylaminonaphthalene-5-Sulphonyl Chloride, *Chemical Industry & Chemical Engineering Quarterly* 16 (1) 31–38.
17. B. Shweta, P. Pares, M. Hiral (2013) Development and Validation of High Performance Thin-Layer Chromatography and Derivative Spectrophotometry methods for determination of Diazepam and Propranolol Hydrochloride in Combined Dosage Form, *International Journal of Drug Development & Research*, 5(1): 91-98.
18. S. Dikran, A. Mohammed, N. Alassaf (2016) Spectrophotometric-Assisted Chemometric Method for the Simultaneous Analysis of Furosemide, Carba-

mazepine, Diazepam, and Carvedilol in their Bulk and Marketed Formulation, *Journal of Natural Science Research*, 6(6): 48-59.

19. S. Dikran, N. Alassaf, F. Zankanah, A. Mohammed (2019) Simultaneous Determination of Furazepam, Carbamazepine, Diazepam and Carvedilol in Quaternary Mixture via Derivative Spectrophotometry, *Journal of Global Pharma Technology*, 11(5) (Suppl.):128-141.

20. V. Sutariya et al.; (2013) Artificial Neural Network in Drug Delivery and Pharmaceutical Research, *The Open Bioinformatics Journal*, 7(1):49-62

21. F. Allegrini and A. C. Olivieri, (2019) *Encyclopedia of Analytical Sciences "Chemometrics and statistics: neural networks"*, Volume 1.

22. V. Kvasnička (1990) An application of neural networks in chemistry, *Chem. Papers*, 44 (6) 775-792

23. G. Goh, N. Hodas, and Abhinav Vishnu (2017) Deep Learning for Computational Chemistry: Review, *Journal of Computational Chemistry* 38: 1291–1307

24. A. Yehia, H. Mohamed (2016) Chemometrics

resolution and quantification power evaluation: Application on Pharmaceutical Quaternary Mixture of Paracetamol, Guaifenesin, Phenylephrine and p-aminophenol, *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 152: 491-500

25. J. Meyer et al.; (2019), Learning Drug Functions from Chemical Structures with Convolutional Neural Networks and Random Forests, *J. Chem. Inf. Model.*, 59: 4438–4449

26. Maltarollo et al.; (2013), Applications of Artificial Neural Networks in Chemical Problems, *InTech*, chapter 10.

27. G. Ioele et al.; (2013), Artificial Neural Network Combined with Principal Component Analysis for Resolution of Complex Pharmaceutical Formulations, *Chem Pharm Bull*, 59(1): 35-40.

28. H. Godini et al.; (2011), Part II: Prediction of the dialysis process performance using Artificial Neural Network (ANN), *Desalination*, 265: 11-21.