

ANALYSIS OF SOME PHENOTHIAZINE DRUGS AND THEIR S-OXIDES USING ENZYMATIC METHOD OF CHOLINESTERASE INHIBITION

ALFRED-UGBENBO D., BLAZHEYEVSKIY M.YE., KOVALENKO V.S., MERZLIKIN S.I.

References:

1. Lally J.; MacCabe, J.H. Antipsychotic medication in schizophrenia: a review. *Br Med Bull.* 2015, 114(1), 169–179. <https://doi.org/10.1093/bmb/ldv017>.
2. Eap, C.B.; Guentert T.W.; Schäublin-Loidl, M.; Stabl, M.; Koeb, L.; Powell, K.; et al. Plasma levels of the enantiomers of thioridazine, thioridazine 2-sulfoxide, thioridazine 2-sulfone, and thioridazine 5-sulfoxide in poor and extensive metabolizers of dextromethorphan and mephenytoin. *Clin Pharmacol Ther.* 1996, 59(3), 322–331. [http://dx.doi.org/10.1016/S0009-9236\(96\)80010-5](http://dx.doi.org/10.1016/S0009-9236(96)80010-5).
3. Kaul, P.N.; Conway, M.W.; Clark, M.L.; Huffine, J. Chlorpromazine metabolism I: Quantitative fluorometric method for 11 chlorpromazine metabolites. *J Pharm Sci.* 1970, 59(12), 1745-1749. <https://doi.org/10.1002/jps.2600591207>.
4. Blazheyevskiy, M.Y. Spectrophotometric and spectrofluorimetric determination of the 2- and 10-disubstituted phenothiazines using peroxy acid oxidation. *Curr. Top. Anal. Chem.* 2019, 11, 67-80. <https://dspace.nuph.edu.ua/handle/123456789/23489?mode=full>.
5. Blazheyevskiy, M.Y.; Dubenska, L.O. An application of derivatization by the peroxic acid oxidation for the determination of fentiasine derivatives by indirect spectrophotometry method (a review). *Odesa Natl Univ Her Chem.* 2019, 24(4(72)), 28–44. [http://dx.doi.org/10.18524/2304-0947.2019.4\(72\).185522](http://dx.doi.org/10.18524/2304-0947.2019.4(72).185522).
6. Daniel, D.; Gutz I.G.R. Spectroelectrochemical determination of chlorpromazine hydrochloride by flow-injection analysis. *J Pharma Biomed Analysis.* 2005, 37(2), 281-286. <https://doi.org/10.1016/j.jpba.2004.10.048>.
7. Kojło, A.; Calatayud, J.M. FIA-spectrophotometric determination of N-substituted phenothiazine derivatives by oxidation with a solid-phase reactor of manganese dioxide incorporated in polyester resin beads. *Talanta.* 1995, 42(7), 909-913. <https://pubmed.ncbi.nlm.nih.gov/18966310/>.
8. Belal, F.; Hadi, H.; Jamal, M. Reversed flow-injection method for estimation of chlorpromazine in pharmaceuticals and urine samples using charge-transfer complexation. *Bull. Chem. Soc. Ethiop.* 2019, 33(1), 11-20. <https://dx.doi.org/10.4314/bcse.v33i1.2>.
9. Chen, D.; Rios, A.; Castro, M.D.; Valcarcel, M. Simultaneous flow-injection determination of chlorpromazine and promethazine by photochemical reaction. *Talanta.* 1991, 38, 1227-1233. [https://doi.org/10.1016/0039-9140\(91\)80097-j](https://doi.org/10.1016/0039-9140(91)80097-j).
10. Sabreen, S.M.; Omar, A.H.; Muna, M.K. Determination of Promethazine Hydrochloride in pharmaceutical forms by Spectrophotometric Method. *Iraqi Academic Scientific Journals.* 2018, 16(3), 28-38. <https://www.iasj.net/iasj/issue/9399>.
11. Puzanowska-Tarasiewicz, H.; Wołyniec, E.; Kojło, A. Flow-injection spectrophotometric determination of promazine. *J. Pharm. Biomed. Anal.* 1996, 14(3), 267 – 271. [https://doi.org/10.1016/0731-7085\(95\)01601-5](https://doi.org/10.1016/0731-7085(95)01601-5).
12. Mohamed, A.M.I.; Abdelmageed, O.H.; Salem, H.; Nagy, D.M.; Omar, M.A. Spectrofluorimetric determination of certain biologically active phenothiazines in commercial dosage forms and human plasma. *Luminescence.* 2013, 28(3), 345-354. <https://doi.org/10.1002/bio.2388>.
13. Mohamed, F.A. Spectrofluorimetric determination of chlorpromazine hydrochloride and thioridazine hydrochloride. *Anal. Lett.* 1995, 28, 2491-2501. <https://doi.org/10.1080/00032719508004030>.
14. Kovalenko, V.S.; Blazheyevskiy, M.Y.; Merzlikin, S.I. A comparative study of the effect of phenothiazine derivatives and their S-oxides on cholinesterase investigated by a new kinetic spectrophotometric method. *Žorg ta farm him.* 2022, 20(1(77)), 35–43. <http://dx.doi.org/10.24959/ophcj.22.252351>.
15. Yang, G.J.; Qu, X.L.; Shen, M.; Qu, Q.S.; Wang, C.Y.; Zhu, A.P.; et al. Trace measurement of phenothiazine drugs in tablets by micellar-enhanced fluorophotometric method. *Fluoresc.* 2007, 17(2), 119–126. <http://doi.org/10.1007/s10895-007-0165-5>.
16. Shetti, P.; Venkatachalam, A. Stability Indicating HPLC Method for Simultaneous Quantification of Trihexyphenidyl Hydrochloride, Trifluoperazine Hydrochloride and Chlorpromazine Hydrochloride from Tablet Formulation. *E-J Chem.* 2010, 7(S1), 299-313. <https://doi.org/10.1155/2010/529386>.
17. Tanaka, E.; Nakamura, T.; Terada, M.; Shinozuka, T.; Hashimoto, C.; Kurihara, K.; et al. Simple and simultaneous determination for 12 phenothiazines in human serum by reversed-phase high-performance liquid chromatography. *Journal of Chromatography.* 2007, 854(1–2), 116–120. <https://doi.org/10.1016/j.jchromb.2007.04.004>.

18. Allender, W.J. High-pressure liquid chromatographic determination of thioridazine and its major metabolites in biological tissues and fluids. *Journal of chromatographic science*. 1985, 23(12), 541-545. <https://doi.org/10.1093/chromsci/23.12.541>.
19. Mehta, A.C. High-performance liquid chromatographic determination of chlorpromazine and thioridazine hydrochlorides in pharmaceutical formulations. *Analyst*. 1981, 106(1267), 1119. <http://dx.doi.org/10.1039/an9810601119>.
20. Geiser, F.; Schultz, M.; Betz, L.; Shaimi, M.; Lee, J.; Champion, W. Direct, preparative enantioselective chromatography of propranolol hydrochloride and thioridazine hydrochloride using carbon dioxide-based mobile phases. *Journal of Chromatography A*. 1999, (1-2), 227-233. [https://doi.org/10.1016/s0021-9673\(99\)01117-6](https://doi.org/10.1016/s0021-9673(99)01117-6).
21. Decaestecker, T.N.; Vande-Castele, S.R.; Wallemacq, P.E.; Van Peteghem, C.H.; Defore, D.L.; Van Bocxlaer, J.F. Information-dependent acquisition-mediated LC-MS/MS screening procedure with semiquantitative potential. *Analytical Chemistry*. 2004, 76(21), 6365-6373. <https://doi.org/10.1021/ac0492315>.
22. Hayen, H.; Karst, U. Analysis of phenothiazine and its derivatives using LC/electrochemistry/MS and LC/electrochemistry/fluorescence. *Analytical Chemistry*. 2003, 75(18), 4833-4840. <https://doi.org/10.1021/ac0346050>.
23. Eap, C.B.; Koeb, L.; Powell, K.; Baumann, P. Determination of the enantiomers of thioridazine, thioridazine 2-sulfone, and of the isomeric pairs of thioridazine 2-sulfoxide and thioridazine 5-sulfoxide in human plasma. *J Chromatogr B. Biomed Appl*. 1995, 669(2), 271-279. [https://doi.org/10.1016/0378-4347\(95\)00096-2](https://doi.org/10.1016/0378-4347(95)00096-2).
24. Asghari, A.; Fahimi, E.; Bazregar, M.; Rajabi, M.; Boutorabi, L. Rapid determination of some psychotropic drugs in complex matrices by tandem dispersive liquid-liquid microextraction followed by high performance liquid chromatography. *J Chromatogr. B. Analyt Technol Biomed Life Sci*. 2017, 1052, 51-59. <https://doi.org/10.1016/j.jchromb.2017.03.012>.
25. Yamini, Y.; Faraji, M. Extraction and determination of trace amounts of chlorpromazine in biological fluids using magnetic solid phase extraction followed by HPLC. *J. Pharm. Anal.* 2014, 4(4), 279-285. <https://doi.org/10.1016/j.jpha.2014.03.003>.
26. Kojlo, A. Indirect potentiometric determination of chlorpromazine with an oxidative column in a flow injection system. *Anal. Lett.* 1997, 30(13), 2353-63. <https://doi.org/10.1080/00032719708001747>.
27. Sales, M.G.; Tomas, J.F.; Lavandeira, S.R. Flow injection potentiometric determination of chlorpromazine. *J. Pharm. Biomed. Anal.* 2006, 41(4), 1280-1286. <http://dx.doi.org/10.1016/j.jpba.2006.03.027>.
28. Daniel, D.; Gutz, I.G. Flow injection spectroelectroanalytical method for the determination of promethazine hydrochloride in pharmaceutical preparations. *Anal. chim. acta*. 2003, 494(1-2), 215-224. [https://doi.org/10.1016/s0003-2670\(03\)00903-6](https://doi.org/10.1016/s0003-2670(03)00903-6).
29. Belal, F.; El-Ashry, S.M.; Shehata, I.M.; El-Sherbeny, M.A.; El-Sherbeny, D.T. Differential-pulse polarographic determination of some N-substituted phenothiazine derivatives in dosage forms and urine through treatment with nitrous acid. *Mikrochim Acta*. 2000, 135(3-4), 147-154. <http://dx.doi.org/10.1007/s006040070004>.
30. Wang, H.; Xu, G.; Dong, S. Electrochemiluminescence sensor using tris(2,2'-bipyridyl) ruthenium(II) immobilized in Eastman-AQ55D-silica composite thin-films. *Anal. Chim. Acta*. 2003, 480(2), 285-290. [https://doi.org/10.1016/s0003-2670\(03\)00049-7](https://doi.org/10.1016/s0003-2670(03)00049-7).
31. Shi, W.B.; Yang, J.D.; Huang, Y.M. Ion-pair complex-based solvent extraction combined with chemiluminescence determination of chlorpromazine hydrochloride with luminol in reverse micelles. *J. Pharm. Biomed. Anal.* 2004, 36(1), 197-203. <https://doi.org/10.1016/j.jpba.2004.05.003>.
32. Mokhtari, A.; Rezaei, B. Chemiluminescence determination of chlorpromazine and fluphenazine in pharmaceuticals and human serum using tris(1,10-phenanthroline) ruthenium(II) *Anal. Methods*. 2011, 3(4), 996-1002. <https://doi.org/10.1039/C0AY00738B>.
33. Hasegawa, C.; Kumazawa, T.; Lee, X.P.; Fujishiro, M.; Kuriki, A.; Marumo, A.; et al. Simultaneous determination of ten antihistamine drugs in human plasma using pipette tip solid-phase extraction and gas chromatography/mass spectrometry. *Rapid Commun Mass Spectrom*. 2006, 20(4), 537-543. <http://dx.doi.org/10.1002/rcm.2335>.
34. Wyjciak-Kosiora, M.; Skalska, A.; Matysik, A. Determination of phenothiazine derivatives by high performance thin-layer chromatography combined with densitometry. *J Pharm Biomed Anal.* 2006, 41(1), 286-289. <https://doi.org/10.1016/j.jpba.2005.10.035>.
35. Trautwein, C.; Kummerer, K. Ready biodegradability of trifluoromethylated phenothiazine drugs, structural elucidation of their aquatic transformation products and identification of environmental risks studied by LC-MSn and QSAR. *Environ Sci Pollut Res*. 2012, 19(8), 3162-3177. <https://doi.org/10.1007/s11356-012-1002-1>.

36. Trautwein, C.; Kummerer, K. Degradation of the tricyclic antipsychotic drug chlorpromazine under environmental conditions, identification of its main aquatic biotic and abiotic transformation products by LC-MSn and their effects on environmental bacteria. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2012, 889:24-38. <https://doi.org/10.1016/j.jchromb.2012.01.022>.
37. Nałęcz-Jawecki, G.; Hajnas, A.; Sawicki, J. Photodegradation and phototoxicity of thioridazine and chlorpromazine evaluated with chemical analysis and aquatic organisms. *Ecotoxicology.* 2008, 17, 13–20. <https://doi.org/10.1007/s10646-007-0171-z>.
38. Wilde, M.L.; Menz, J.; Trautwein, C.; Leder, C.; Kümmerer, K. Environmental fate and effect assessment of thioridazine and its transformation products formed by photodegradation. *Environ Pollut.* 2016, 213, 658–670. <http://dx.doi.org/10.1016/j.envpol.2016.03.018>.
39. Castillo-Zacarías, C.; Barocio, M.E.; Hidalgo-Vázquez, E.; et al., Antidepressant drugs as emerging contaminants: Occurrence in urban and non-urban waters and analytical methods for their detection. *Sci Total Environ.* 2021, 757, 143722. doi: 10.1016/j.scitotenv.2020.143722.
40. Radic, Z.; Pickering, N.A.; Vellom DC.; Camp, S.; Taylor, P. Three distinct domains in the cholinesterase molecule confer selectivity for acetyl and butyrylcholinesterase inhibitors. *Biochemistry.* 1993, 32(45), 12074-12084. <https://doi.org/10.1021/bi00096a018>.
41. Saxena, A.; Redman, A.M.; Jiang, X.; Lockridge, O.; Doctor, B.P. Differences in active site gorge dimensions of cholinesterases revealed by binding of inhibitors to human butyrylcholinesterase. *Biochemistry.* 1997, 36(48), 14642–14651. <http://dx.doi.org/10.1021/bi971425>
42. Sanajou, S.; Nourhashemi, S.; Fallah, A.; Ercetin, T.; Sahin, M.F.; Gulcan, H.O. The investigation of the interaction of several antipsychotic drugs with human cholinesterase enzymes. *EMUJPharmSci.* 2018, 1(1), 1-5. <https://ojs.emu.edu.tr/index.php/jps/article/view/27>.
43. Debord, J.; Merle, L.; Bollinger J-C.; Dantoine, T. Inhibition of butyrylcholinesterase by phenothiazine derivatives. *J Enzyme Inhib Med Chem.* 2002, 17(3), 197–202. <http://dx.doi.org/10.1080/1475636021000003165>.
44. De los Ríos C. Cholinesterase inhibitors: a patent review (2007 – 2011). *Expert Opin Ther Pat.* 2012, 22(8), 853–69. <http://dx.doi.org/10.1517/13543776.2012.701619>.
45. Darvesh, S.; McDonald R.S.; Penwell, A.; Conrad, S.; Darvesh, K.V.; Mataija, D.; et al. Structure-activity relationships for inhibition of human cholinesterases by alkyl amide phenothiazine derivatives. *Bioorg Med Chem.* 2005, 13(1), 211–222. <http://dx.doi.org/10.1016/j.bmc.2004.09.059>.
46. Darvesh, S.; Macdonald, I.R.; Martin, E. Selectivity of phenothiazine cholinesterase inhibitors for neurotransmitter systems. *Bioorg Med Chem Lett.* 2013, 23(13), 3822–3825. <http://dx.doi.org/10.1016/j.bmcl.2013.04.082>.
47. Darvesh, S.; Pottie, I.R.; Darvesh, K.V.; McDonald, R.S.; Walsh, R.; Conrad, S.; et al. Differential binding of phenothiazine urea derivatives to wild-type human cholinesterases and butyrylcholinesterase mutants. *Bioorg Med Chem.* 2010, 18(6), 2232–2244. <http://dx.doi.org/10.1016/j.bmc.2010.01.066>.
48. Evtugin, G.A.; Budnikov, G.K.; Nikolskaya, E.B. Biochemical Tests Based on Stabilized Cholinesterase Preparations. *New Approaches. J Anal Chem.* 2002, 57(10), 954–959. <http://dx.doi.org/10.1023/a:1020491428970>.
49. Blazheyevskiy, M.Ye.; Dyadchenko, V.V. Kinetic determination of cholinesterase inhibitors by biochemical method using the oxidation reaction of p-phenetidine as an indicator. *Farmatsevtichnyi zhurnal.* 2004, 2, 52-58. <https://pharmj.org.ua/archives/2004/2004-2.pdf>
50. Parker, W.E.; Witnauer, L.P.; Swern, D. Peroxides. IV.2 aliphatic diperoxides. *J Am Chem Soc.* 1957, 79(8), 1929–1931. <http://dx.doi.org/10.1021/ja01565a043>.
51. Kovalenko, V. S.; Blazheyevskiy, M. Ye.; Merzlikin, S. I. A comparative study of the effect of phenothiazine derivatives and their S-oxides on cholinesterase investigated by a new kinetic spectrophotometric method. *J. Org. Pharm. Chem.* 2022, 20(1), 35-43. <https://doi.org/10.24959/ophcj.22.252351>