

ANALYTICAL VALIDATION STUDY WITH UNCERTAINTY EVALUATION OF THE METHOD FOR QUANTITATIVE DETERMINATION OF HESPERIDIN IN THE EXTRACTED PRODUCT AND MANDARIN PROCESSING WASTE

IMEDA RUBASHVILI, MZIA TSITSAGI, MARIAM CHKHAIDZE, KETEVAN EBRALIDZE

References:

1. Ellouze, I. Citrus bio-wastes: a source of bioactive, functional products and non-food uses. In: Mediterranean fruits biowastes. 2022, 221–260. https://doi.org/10.1007/978-3-030-84436-3_9
2. Samota, M.K.; Kaur, M.; Sharma, M.; Sarita, Krishnan, V.; Thakur, J.; Rawat, M.; Phogat, B.; P.N. Guru. Hesperidin from citrus peel waste: extraction and its health implications. *Qual. Assur. Saf. Crops Foods*. 2023, 15(2), 71–99. <https://doi.org/10.15586/qas.v15i2.1256>
3. Dukare, A.; Bibwe, B.; Samota, M.K.; Dawange, S.; Kumar, M.; Lorenzo, J.M.; Assessment of bioactive compounds, physicochemical properties and microbial attributes of hot air-dried mango seed kernel powder: an approach for quality and safety evaluation of hot air-dried mango seed kernel powder. *Food Anal. Methods*. 2022, 15(10), 2675–2690. <https://doi.org/10.21203/rs.3.rs-1631292/v1>
4. Aggarwal, V.; Tuli, H.; Thakral, F.; Singhal, P.; Aggarwal, D.; Srivastava, S.; Pandey A.; Sak, k.; Varol, Md., M.; Khan, A., Md.; Sethi, G. Molecular mechanisms of action of hesperidin in cancer: recent trends and advancements. *Exp. Biol. Med*. 2020, 245(5), 486–497. <https://doi.org/0.1177/1535370220903671>
5. Pyrzynska, K.; Hesperidin: A Review on Extraction Methods, Stability and Biological Activities. *Nutrients* 2022, 14(2), 2387. <https://doi.org/10.3390/nu14122387>
6. Rubashvili, I.; Tsitsagi, M.; Zautashvili, M.; Chkhaidze, M.; Ebralidze, K.; Tsitsishvili, V. Extraction and analysis of oleanolic acid and ursolic acid from apple processing waste materials using ultrasound-assisted extraction technique combined with high performance liquid chromatography. *Rev. Roum. Chim*. 2020, 65(10), 919-928. <https://doi.org/10.33224/rrch.2020.65.10.07>
7. Ayinzat, F.; Jo, A.; Bawa, E.K. Hesperidin-sources, chemistry, extraction, measurement and biologic effects on reproduction in animals: a review. *Int. J. Vet. Sci. Anim. Husb*. 2021, 6(4), 1–8. <https://doi.org/10.22271/veterinary.2021.v6.i3a.360>
8. Boudhrioua, M. N.; M'hiri, N.; Ioannou, I.; Paris, C.; Ghoul, M. Comparison of the efficiency of different extraction methods on antioxidants of maltase orange peel. *Int. J. Food. Sci. Nutr*. 2016, 3(2), 1–13. <https://doi.org/10.15436/2377-0619.16.789>
9. M'hiri, N.; Ioannou, I.; Boudhrioua, N.M. Ghoul, M. Comparison of the efficiency of different extraction methods on antioxidants of maltease orange peel. *Int. J. Food. Nutr. Sci*. 2015, 3(1), 239–251. <https://doi.org/10.15436/2377-0619.16.789>
10. M'hiri, N.; Ioannou, I.; Mihoubi Boudhrioua, N. Ghoul, M. Effect of different operating conditions on the extraction of phenolic compounds in orange peel. *Food Bioprod. Process*. 2015, 96, 161–170. <https://doi.org/10.1016/j.fbp.2015.07.010>
11. Sharma, P.; Pandey, P.; Gupta, R.; Roshan, S.; Garg, A.; Shulka A.; Pasi, A. Isolation and characterization hesperidin from orange peel. *Indo Am. J. Pharm. Sci*. 2013, 3(4), 3892–3897
12. M'hiri, N.; Ioannou, I.; Ghoul, M.; Boudhrioua, N. Extraction methods of citrus peel phenolic compounds. *Food Rev. Int*. 2014, 30(4), 265–290. <https://doi.org/10.1080/87559129.2014.924139>
13. Hayat, K.; Hussain, S.; Abbas, S.; Farooq, U.; Ding, B.; Xia, S.; Jia, C.; Zhang, X.; Xia, W. Optimized microwave-assisted extraction of phenolic acids from citrus mandarin peels and evaluation of antioxidant activity in vitro. *Sep. Purif. Technol*. 2009, 70(1), 63–70. <https://doi.org/10.1016/j.seppur.2009.08.012>
14. Khan, M.; Abert-Vian, M.; Fabiano-Tixier, A.; Dangles, O.; Chemat, F. Ultrasound-assisted extraction of polyphenols (flavanone glycosides) from orange (*Citrus sinensis* L.) peel. *Food Chemistry*. 2010, 119(2), 851–858. <https://doi.org/10.1016/j.foodchem.2009.08.046>
15. Li, B.B.; Smith, B; Hossain, M.M.; Extraction of phenolics from citrus peels: I. Solvent extraction method. *Sep. Purif. Technol*. 2006, 48(2), 182–188. <https://doi.org/doi.org/10.1016/j.seppur.2005.07.005>
16. Li, C.; Schluesener, H. Health-promoting effects of the citrus flavanone hesperidin. *Critical Reviews in Food Sci. Nutr*. 2016, 57(3), 613–631. <https://doi.org/10.1080/10408398.2014.906382>
17. Tsitsagi, M.; Rubashvili, I.; Chkhaidze, M.; Khachidze, M.; Ebralidze, K. Four Valuable Bioactive Compounds from Citrus Waste Using Sequential Two-Step Ultrasound-Assisted Extraction Method. *Bull. Georgian Natl. Acad. Sci*. 2021, 15(4), 50-56
18. Tsitsagi, M.; Ebralidze, K.; Chkhaidze, M.; Rubashvili, I.; Tsitsishvili, V. Sequential Extraction of Bioactive Compounds from Tangerine (*Citrus Unshiu*) Peel. *Ann. Agrar. Sci*. 2018, 16(2), 236-241. <https://doi.org/10.1016/j.aasci.2018.02.007>

19. Saeidi, I.; Hadjmohammadi, M.R.; Peyrovi, M.; Iranshahi, M.; Barfi, B.; Babaei, A.B.; Dust, A.M. HPLC determination of hesperidin, diosmin and eriocitrin in Iranian lime juice using polyamide as an adsorbent for solid phase extraction. *J. Pharm. Biomed. Anal.* 2011, 56(2), 419-422. <https://doi.org/10.1016/j.jpba.2011.05.015>
20. Kanaze, F.I.; Gabrieli, C.; Kokkalou, E.; Georgarakis, M.; Niopas, I. Simultaneous reversed-phase high-performance liquid chromatographic method for the determination of diosmin, hesperidin and naringin in different citrus fruit juices and pharmaceutical formulations. *J. Pharm. Biomed. Anal.* 2003, 33(2), 243–2499. [https://doi.org/10.1016/s0731-7085\(03\)00289-9](https://doi.org/10.1016/s0731-7085(03)00289-9)
21. Tsai, T.H.; Liu, M.C. Determination of extracellular hesperidin in blood and bile of anaesthetized rats by microdialysis with high performance liquid chromatography, a pharmacokinetic application. *J. Chromatogr. B. Analyt. Technol. Biomed. Life. Sci.* 2004, 806(2), 161–166. <https://doi.org/10.1016/j.jchromb.2004.03.047>
22. Kanaze, F.I.; Kokkalou, E.; Georgarakis, M.; Niopas, I. A validated solid-phase extraction HPLC method for the simultaneous determination of the citrus flavanone aglycones hesperidin and naringenin in urine. *J. Pharm. Biomed. Anal.* 2004, 36(1), 175–181. <https://doi.org/10.1016/j.jpba.2004.05.015>
23. Adham, A.N. Qualitative and Quantitative Estimation of Hesperidin in Peel and Juice of Citrus Fruits by RP-HPLC Method Growing in Kurdistan Region/Iraq. *Int. J. Pharm. Sci. Rev. Res.* 2015, 33(2), 220-224.
24. Chen, L.; Zhu, S.; Wang, C.; Chen, L. Development of a HPLC-UV method for the separation and quantification of hesperidin, neohesperidin, neohesperidin dihydrochalcone and hesperetin. *Nat. Prod.* 2023, 37(10), 1714-1718. <https://doi.org/10.1080/14786419.2022.2104275>
25. International Conference on Harmonization, Harmonized Tripartite guideline, Validation of analytical procedures, text and methodology Q2 (R1). 2005
26. Rubashvili, I.; Tsitsagi, M.; Ebralidze, K.; Tsitsishvili, V.; Eprikashvili, L. Chkhaidze, M.; Zautashvili, M. Extraction and Analysis of the Major Carotenoids of Agro-Industrial Waste Materials using Sequential Extraction Techniques and High-Performance Liquid Chromatography. *Eurasian J. Anal. Chem.* 2018, 13(2), em06
27. Das, A.K.; Dewanjee, S. Chapter 3 - Optimization of Extraction Using Mathematical Models and Computation: in Computational Phytochemistry. *Computational Phytochemistry*. 2018, 75-106
28. PA/PH/OMCL (18) 145 R1 CORR. Evaluation of Measurement Uncertainty (Core Document). 2020
29. Ellison, S.L.R.; Rosslein, M.; Williams, A. Quantifying uncertainty in analytical measurement. EURACHEM/ CITAG Guide CG4. Third edition. Eurachem/CITAC. 2012
30. EA-4/16 G:2003. EA guidelines on the expression of uncertainty in quantitative testing. 2003
31. Rubashvili, I.; Tsitsagi, M.; Ebralidze, K. Measurement uncertainty evaluation of analytical method for quantitative determination of ursolic and oleanolic acids in apple processing agroindustrial waste material. *Per. Tchê Quim.* 2022, 19(42), 60-76. https://doi.org/10.52571/PTQ.v19.n42.2022_06_RUBASH-VILI_pgs_60_74.pdf
32. Inoue, T.; Tsubaki, S.; Ogawa, K.; Onishi, K.; Azuma, J. Isolation of hesperidin from peels of thinned Citrus unshiu fruits by microwave-assisted extraction. *Food Chem.* 2010, 123(2), 542-547. <https://doi.org/10.1016/j.foodchem.2010.04.051>
33. Šafranko, S.; Corkovic, I.; Jerkovic, I.; Jakovljevic, M.; Aladic, K.; Šubaric, D.; Jokic, A.; Green extraction techniques for obtaining bioactive compounds from mandarin peel (Citrus unshiu var. Kuno): Phytochemical Analysis and Process Optimization. *Foods* 2021, 10(5), 1043. <https://doi.org/10.3390/foods10051043>
34. Papoutsis, K.; Pristijono, P.; Golding, J.B.; Stathopoulos, C.E.; Bowyer, M.C.; Scarlett, C.J.; Vuong, Q. V. Screening the effect of four ultrasound-assisted extraction parameters on hesperidin and phenolic acid content of aqueous citrus pomace extracts. *Food Biosci.* 2018, 21, 20-26. <https://doi.org/10.1016/j.fbio.2017.11.001>
35. Gómez-Mejía, E.; Rosales-Conrado, N.; León-González M.E.; Madrid Y. Citrus peels waste as a source of value-added compounds: Extraction and quantification of bioactive polyphenols. *Food Chem.* 2019, 295, 15, 289-299. <https://doi.org/10.1016/j.foodchem.2019.05.136>
36. Suri, S.; Singh, A.; Nema, P.K. Current applications of citrus fruit processing waste: A scientific outlook. *Applied Food Research.* 2022, 2(1), 100050. <https://doi.org/10.1016/j.afres.2022.100050>