

GREEN CHEMISTRY-BASED INVESTIGATION OF PASSIFLORA EXTRACTS: ANTIOXIDANT AND ANTICHOLINESTERASE ACTIVITIES TARGETING ALZHEIMER'S DISEASE MECHANISMS

GİZEM İĞDELİ, DEMET DİNCEL, YASİN DARI, CAGLA KIZILARSLAN HANCER, MURAT KARTAL, GULACTI TOPCU

References:

1. Amoateng P, et al. Medicinal plants used in the treatment of mental and neurological disorders in Ghana. *Evid Based Complement Alternat Med*. 2018;2018:8590381.
2. Mostefa N, et al. Anti-Alzheimer's activity of polyphenolic stilbene-rich acetone fraction of the oil-removed seeds of *Passiflora edulis*: in vivo and in silico studies. *Chem Biodivers*. 2023;20(5):e202201051.
3. Ingale SP, Kasture SB. Protective effect of standardized extract of *Passiflora incarnata* flower in Parkinson's and Alzheimer's disease. *Anc Sci Life*. 2017;36(4):200-6.
4. Cao SQ, et al. *P. edulis* extract protects against amyloid- β toxicity in Alzheimer's disease models through maintenance of mitochondrial homeostasis via the FOXO3/DAF-16 pathway. *Mol Neurobiol*. 2022;59(9):5612-29.
5. Ingale SP, Joshi AM, Ingale PL. Network pharmacology-based evaluation of therapeutic potential of *Passiflora edulis* leaves in Alzheimer's disease. *Adv Pharmacol Pharm*. 2024;12(3):186-98.
6. Kim GH, et al. Improvement in neurogenesis and memory function by administration of *Passiflora incarnata* L. extract applied to sleep disorder in rodent models. *J Chem Neuroanat*. 2019;98:27-40.
7. Akhondzadeh S, et al. Passionflower in the treatment of generalized anxiety: A pilot double-blind randomized controlled trial with oxazepam. *J Clin Pharm Ther*. 2001;26(5):363-7.
8. Jawna-Zbońska K, et al. *Passiflora incarnata* L. improves spatial memory, reduces stress, and affects neurotransmission in rats. *Phytother Res*. 2016;30(5):781-9.
9. Sarris J, et al. Herbal medicine for depression, anxiety, and insomnia: a review of psychopharmacology and clinical evidence. *Eur Neuropsychopharmacol*. 2011;21(12):841-60.
10. Trotti LM, Karroum EG. Melatonin for sleep disorders in patients with neurodegenerative diseases. *Curr Neurol Neurosci Rep*. 2016;16:1-10.
11. Schmidt BM, Cheng DM. *Ethnobotany: A phytochemical perspective*. John Wiley & Sons; 2017.
12. Demirezer LÖ, Ersöz T, Saraçoğlu İ, Şener B. *Tedavide Kullanılan Bitkiler FFD Monografıları*. Nobel Tıp Kitabevleri; 2007.
13. Abbasi E, et al. Neuroprotective effects of vitexin, a flavonoid, on pentylene-tetrazole-induced seizure in rats. *Chem Biol Drug Des*. 2012;80(2):274-8.
14. Choi JS, et al. Effects of C-glycosylation on anti-diabetic, anti-Alzheimer's disease and anti-inflammatory potential of apigenin. *Food Chem Toxicol*. 2014;64:27-33.
15. Malar DS, et al. Vitexin inhibits A β 25-35 induced toxicity in Neuro-2a cells by augmenting Nrf-2/HO-1 dependent antioxidant pathway and regulating lipid homeostasis by the activation of LXR- α . *Toxicol In Vitro*. 2018;50:160-71.
16. Scullin MK, Bliwise DL. Sleep, cognition, and normal aging: integrating a half century of multidisciplinary research. *Perspect Psychol Sci*. 2015;10(1):97-137.
17. Villa C, Ferini-Strambi L, Combi R. The synergistic relationship between Alzheimer's disease and sleep disorders: an update. *J Alzheimers Dis*. 2015;46(3):571-80.
18. Ellman GL, et al. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol*. 1961;7(2):88-95.
19. Kızıltaş H, et al. LC-HRMS profiling and antidiabetic, anticholinergic, and antioxidant activities of aerial parts of kınkor (*Ferulago stellata*). *Molecules*. 2021;26(9):2469.
20. Bingol Z, et al. Antidiabetic, anticholinergic and antioxidant activities of aerial parts of shaggy bindweed (*Convolvulus betonicifolia* Miller subsp.)—profiling of phenolic compounds by LC-HRMS. *Heliyon*. 2021;7(5):eXXXXXX.
21. Eruçar S. Bazı bitkisel çayların fenolik madde profili ve antioksidan aktivitelerinin incelenmesi. *Fen Bilimleri Enstitüsü*; 2006.
22. Slinkard K, Singleton VL. Total phenol analysis: automation and comparison with manual methods. *Am J Enol Vitic*. 1977;28(1):49-55.
23. Moreno MİN, et al. Comparison of the free radical-scavenging activity of propolis from several regions of Argentina. *J Ethnopharmacol*. 2000;71(1-2):109-14.
24. Komersova A, Komers K, Čegan A. New findings about Ellman's method to determine cholinesterase activity. *Z Naturforsch C*. 2007;62(1-2):150-4.

25. Wojnowski W, et al. AGREEprep—analytical greenness metric for sample preparation. *TrAC Trends Anal Chem.* 2022;149:116553.
26. Aman U, et al. *Passiflora incarnata* attenuation of neuropathic allodynia and vulvodinia apropos GABA-ergic and opioidergic antinociceptive and behavioural mechanisms. *BMC Complement Altern Med.* 2016;16:1-17.
27. Waszkowiak K, Gliszczyńska-Świgło A. Binary ethanol–water solvents affect phenolic profile and antioxidant capacity of flaxseed extracts. *Eur Food Res Technol.* 2016;242:777-86.
28. Chaabani E, et al. Ethanol–water binary solvent affects phenolic composition and antioxidant ability of *Pistacia lentiscus* L. fruit extracts: a theoretical versus experimental solubility study. *J Food Meas Charact.* 2023;17(5):4705-14.
29. Ziani I, et al. The effect of ethanol/water concentration on phenolic composition, antioxidant, and antimicrobial activities of *Rosmarinus tournefortii* de Noé hydrodistillation solid residues. *J Food Meas Charact.* 2023;17(2):1602-15.