

FLUORIMETRIC DERIVATIZATION-BASED HPLC-FL METHOD FOR THE PROTOTYPE PHARMACOKINETIC ANALYSIS OF SELEXIPAG IN HUMAN PLASMA

BURHAN CEYLAN, MELTEM ÇAYCI, CEM ÖNAL, ARMAĞAN ÖNAL

References:

1. Richter, M. J.; Gall, H.; Grimminger, J.; Grimminger, F.; Ghofrani, H-A. Selexipag for the treatment of pulmonary arterial hypertension. *Expert Opin Pharmacother.* 2016, 17 (13), 1825-1834. <https://doi.org/10.1080/14656566.2016.1215429>
2. Simonneau, G.; Torbicki, A.; Hoeper, M. M.; et al. Selexipag: an oral, selective prostacyclin receptor agonist for the treatment of pulmonary arterial hypertension. *Eur. Respir. J.* 2012, 40 (4), 874-880. <https://doi.org/10.1183/09031936.00137511>
3. Kaufmann, P.; Hurst, N.; Astruc, B., Dingemanse, J. Absolute oral bioavailability of selexipag, a novel oral prostacyclin IP receptor agonist. *Eur. J. Clin. Pharmacol.* 2017, 73 (2), 151-156. <https://doi.org/10.1007/s00228-016-2164-4>
4. Gorumutçu, G. P.; Ratnakaram, V. N. Oxidative Coupling: A tranquil approach for determination of selexipag by visible spectrophotometry. *Orient. J. Chem.* 34 (6), 3112-3117. <http://dx.doi.org/10.13005/ojc/340656>
5. Li, L.; Wang, H.; Tian H.; Li, J., et al. Determination of 6 residual solvents in selexipag by headspace gas chromatography. *J. China Pharma. Univ.* 2018, 49 (1), 93-96. <http://dx.doi.org/10.11665/j.issn.1000-5048.20180113>
6. Santosh, S. S.; Devi, N.K. Stability indicating method development and validation of selexipag in bulk and pharmaceutical dosage form by using RP-HPLC. *J. Pharm. Res.*, 2021, 5 (2), 000239. <https://doi.org/10.23880/oajpr-16000239>
7. Xie, S.; Shi, L.; Chen, J.; et al. Simultaneous quantification and pharmacokinetic investigation of selexipag and its main metabolite ACT-333679 in rat plasma by UPLC-MS/MS method. *J. Pharm. and Biomed. Anal.* 2020, 190, 113496. <https://doi.org/10.1016/j.jpba.2020.113496>
8. Rao, K. P.; Babu, N. L.; Koganti, K.; et al. Related substances method development and validation of an LC-MS/MS method for quantification of selexipag and its related impurities in rat plasma and application to pharmacokinetic studies. *SN App. Sci.* 2021, 3, 3, 321. <https://link.springer.com/article/10.1007/s42452-021-04219-x>
9. Ceylan, B.; Tiris, G.; Tekkeli Kepekci, S. E.; et al. A novel HPLC method for selexipag in human plasma and application to a prototype pharmacokinetic study. *Acta Chrom.* 2024, 36, 3, 273-278. <https://doi.org/10.1556/1326.2023.01167>
10. Satheshkumar, V.; Muruganantham, V. A rabbit method for quantification of selexipag in human plasma using high performance liquid chromatography with electron spray ionization tandem mass spectrometry. *Annals R.S.C.B.* 2021, 25, 4, 5689-5707.
11. Fager, R. S.; Kutina, C. B.; Abrahamson, E. W. The use of NBD chloride (7 chloro-4-nitro-2-oxa-1,3-diazole) in detecting amino acids and as an N-terminal reagent. *Anal Biochem.* 1973, 53, 1, 290-294. [https://doi.org/10.1016/0003-2697\(73\)90433-8](https://doi.org/10.1016/0003-2697(73)90433-8)
12. Tekkeli Kepekci, S. E.; Önal, A. Spectrofluorimetric methods for the determination of gemifloxacin in tablets and spiked plasma samples. *J. Fluoresc.* 2011, 21, 1001-1007. <https://doi.org/10.1007/s10895-010-0759-1>
13. Job, P. Study of new selective reagent acetophenone 2',4'- Dihydroxy semicarbazone for extractive spectrophotometric determination of vanadium. *Ann. Chim.* 1928, 9, 113-203.
14. EMA (2011). European Medicines Agency (EMA)/Committee for Medicinal Products for Human Use (CHMP). Guideline on Bioanalytical Method Validation, EMA/CHMP, London, United Kingdom.
15. Tiris, G.; Tekkeli Kepekci S. E.; Önal, C.; et al. An UPLC method for the determination of sorafenib in human plasma by fluorimetric detection with pre-column derivatization and application to a pharmacokinetic study. *Rev. Roum. Chim.* 2022, 67, 3, 217-224. <https://doi.org/10.33224/rch.2022.67.3.10>
16. G. Heinzel, R. Woloszczak, S. Thomann, Version TOPFITT 2.0 Pharmacokinetic and Pharmacodynamic Data Analysis System for the PC. VCH Publishers: Gustav-Fisher-Verlag, Stuttgart, Germany, (1993).