

DETERMINATION OF GUAIFENESIN IN SPIKED HUMAN BREAST MILK: HPLC-UV METHOD DEVELOPMENT, VALIDATION, AND UNCERTAINTY EVALUATION

BURHAN CEYLAN, CEM ÖNAL, ARMAĞAN ÖNAL

References:

1. Green, J. L.; Wang, G. S.; Reynolds, K. M.; et al. Safety profile of cough and cold medication use in paediatrics. *Pediatrics* 2017; 139 (6), e20163070. <https://doi.org/10.1542/peds.2016-3070>
2. Kuhn, J. J.; Hendley, J. O.; Adams, K. F.; Clark, J. W.; Gwaltney, J. M. Antitussive effect of guaifenesin in young adults with natural colds. Objective and subjective assessment. *Chest* 1982; 82 (6), 713-718. <https://doi.org/10.1378/chest.82.6.713>
3. Dicpinigaitis, P. V.; Gale, Y. E. Effect of guaifenesin on cough reflex sensitivity. *Chest* 2003; 124 (6), 2178-2181. <https://doi.org/10.1378/chest.124.6.2178>
4. Aluri, J. B.; Stavchansky, S. Determination of guaifenesin in human plasma by liquid chromatography in the presence of pseudoephedrine. *J. Pharmaceu. Biomed. Analy.* 1993; 11 (9), 803-808. [https://doi.org/10.1016/0731-7085\(93\)80072-9](https://doi.org/10.1016/0731-7085(93)80072-9)
5. Kulikov, A. U.; Verushkin, A. G. Simultaneous determination of paracetamol, caffeine, guaifenesin and preservatives in syrups by micellar LC. *Chromatographia* 2008; 67, 347-355. <https://doi.org/10.1365/s10337-007-0510-5>
6. Hatami, M.; Farhadi, K.; Abdollahpour, A. Using dispersive liquid-liquid microextraction and liquid chromatography for determination of guaifenesin enantiomers in human urine. *J. Sep. Sci.* 2011; 34, 2933-2939. <https://doi.org/10.1002/jssc.201100520>
7. El-Gindy, A.; Emara, S.; Shaaban, H. Development and validation of chemometrics-assisted spectrophotometric and liquid chromatographic methods for the simultaneous determination of two multicomponent mixtures containing bronchodilator drugs. *J. Pharmce. Biomed. Analy.* 2007; 43 (3), 973-982. <https://doi.org/10.1016/j.jpba.2006.09.020>
8. Bankar, A. A.; Lokhande, S. R.; Sawant, R. L.; Bhagat, A. R. Spectrophotometric estimation of guaifenesin and salbutamol in pure and tablet dosage form by using different methods. *Der. Pharma. Chemica* 2013; 5 (3), 92-97.
9. Kamatham, S. S.; Kolli, S.; Joga, D. D.; Vasa, K. R.; Yanamadala, B. D. Simultaneous estimation of salbutamol, ambroxol and guaifenesin in tablet dosage forms by using RP-HPLC. *Int. J. Pharm. Res. Biomed. Analy.* 2013; 2 (3), 1-13.
10. Walode, S. G.; Deshpande, S. D.; Deshpande, A. V. Stability indicating RP-HPLC method for simultaneous estimation of salbutamol sulphate and guaifenesin. *Pharmacia Sinica* 2013; 4 (2), 61-67.
11. Ezhava, S. B.; Bhalara, K. D.; Rathod, I. S.; Bhalara, D. D. Simultaneous estimation of salbutamol sulphate and guaifenesin in their combined liquid dosage form by HPTLC method. *Int. J. Pharm. Res. Healt. Sci.* 2014; 2 (2), 191-196.
12. Wen, J.; Zhang, H.; Xia, C.; Hu, X.; et al. A sensitive liquid chromatography-electrospray ionization-mass spectrometry method for the simultaneous determination of pentoxifyverine citrate and guaifenesin in human plasma-application to pharmacokinetic and bioequivalence studies. *Biomed. Chrom.* 2010; 24, 351-357. <https://doi.org/10.1002/bmc.1298>
13. Maresova, V.; Chadt, J.; Novakova, E. Screening and semiquantitative analysis of drugs and drugs of abuse in human serum samples using gas chromatography-mass spectrometry. *Neuroendocrinology Letters* 2008; 29 (5), 749-754.
14. Yehia, A. M.; Mohamed, H. M. Chemometrics resolution and quantification power evaluation: application on pharmaceutical quaternary mixture of paracetamol, guaifenesin, phenylephrine and p-aminophenol. *Spectrochimica Acta Part A-Molecular and Biomolecular Spectroscopy* 2016; 152, 491-500.
15. Abdelwahab, N. S.; Abdelaleem, E. A. TLC-densitometric determination of guaifenesin, pseudoephedrine hydrochloride and guaifenesin related substance (Guaiacol). *Journal of Planar Chromatography* 2013; 26, 73-77.
16. EMA (2011). European Medicines Agency (EMA)/Committee for Medicinal Products for Human Use (CHMP). Guideline on Bioanalytical Method Validation, EMA/CHMP, London, United Kingdom.