

DETERMINATION OF QUERCETIN AND SUM OF POLYPHENOLS BY USING 18-MOLYBDODIPHOSPHATE HETEROPOLY COMPLEX

T.A. DENISENKO, L.P. TSIGANOK, A.B. VISHNIKIN. SPECTROPHOTOMETRIC

References:

1. Brett A.M.O., Ghica M.-E. Electrochemical oxidation of quercetin. *Electroanalysis*. 2003, 15 (22), 1745-1750.
2. Tarakhovskii Yu. S., Kim Yu. A., Abdrasilov B. S., Muzafarov E. N. Flavonoidy: biokhimiya, biofizika, meditsina. Pod red. E.I. Maevskii. Pushchino: Sunchrobook, 2013. S. 310.
3. Rastitel'nye lekarstvennye sredstva. Pod red. N.P. Maksyutinoi. Kiev, 1985, S. 285.
4. Khimicheskii analiz lekarstvennykh rastenii. Pod redaktsiei Grinkevich H.I., Safronich L. H. Moskva: Vysshaya shkola, 1983. S. 176.
5. Biesaga M., Pyrzynska K. Analytical procedures for determination of quercetin and its glycosides in plant material. *Crit. Rev. Anal. Chem.* 2009, 39, 95-107
6. Kurkina A.V., Ryzhov V.M., Avdeeva E.V. Perspektivy ispol'zovaniya vysokoeffektivnoi zhidkostnoi khromatografii dlya standartizatsii syr'ya i preparatov bessmertnika peschanogo. *Izvestiya Samarskogo nauchnogo tsentra Rossiiskoi akademii nauk*. 2011, 13 (1/8), 144-150.
7. Khodakov I.V. Sposob identifikatsii polifenolov v rastitel'nykh ekstraktakh pri pomoshchi VEZhKh. *Opreделение состава изофлавонов сои. Методы и объекты хим. анализа*. 2013, 8(3), 132-142.
8. Geng C. H., Lin M., Wang W. Y., Ye J. N.. *Opreделение aktivnykh ingredientov v boyaryshnike i ego komponentakh metodom kapillyarnogo elektroforeza s elektrokhimicheskimi detektirovaniem*. *Zhurn. analit. khimii*. 2008, 63(1), 455-462.
9. Ignat I., Volf I., Popa V.I. A critical review of methods for characterisation of polyphenolic compounds in fruits and vegetables. *Food Chem.* 2011, 126, 1821-1835.
10. Hassan H.N.A., Barsoum B.N., Habib I.H.I. Simultaneous spectrophotometric determination of rutin, quercetin and ascorbic acid in drugs using a Kalman Filter approach. *J. Pharm. Biomed. Anal.* 1999, 20, 315-320.
11. Giusti, M.M., Wrolstad R.E. Acylated anthocyanins from edible sources and their applications in food systems. *Biochemical Engineering Journal*. 2003, 14(3), 217-225.
12. Zaitsev V.M., Khalaf V.A., Zaitseva G.M. Metodi kontsentruvaniya ta viznachennya fenol'nikh spoluk. *Metody i ob'ekty khim. analiza*. 2008, 3(1), 4-21.
13. 4-21.
14. Korenman I.M. Fotometricheskii analiz. *Metody opredeleniya organicheskikh soedinenii*. Moskva: Khimiya, 1975. S. 359.
15. Lobanova A.A., Budaeva V.V., Sakovich G.V. Issledovanie biologicheskii aktivnykh flavonoidov v ekstraktakh iz rastitel'nogo syr'ya. *Khimiya rastitel'nogo syr'ya*. 2004, 1, 47-52.
16. Box, J. D. Investigation of the Folin-Ciocalteu phenol reagent for the determination of polyphenolic substances in natural waters. *Water Res.* 1983, 17, 511-525.
17. Singleton V.L., Orthofer R., Lamuela-Raventos R.M. Analysis of total phenols and other oxidations substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in enzymology*. 1999, 299, 152-178.
18. Bhatia I.S., Singh J., Bajaj K.L. A sensitive colorimetric method for the microdetermination of flavonols. *Microchim. Acta*. 1974, 62(5), 909-913.
19. Sakamoto M., Takamura K. Consecutive determination of rutin and quercetin by spectrophotometric measurements. *Microchem. J.* 1978, 23, 374-381.
20. Dowd L.E. Spectrophotometric determination of quercetin. *Anal. Chem.* 1959, 31(7), 1184-1187.
21. Jurd L. Aluminium complexes of phenolic flavones. Spectral and structural correlation. *Phytochemistry*. 1969, 8, 445-462.

22. Kaushal G.P., Sekhon B.S., Bhatia I.S. Spectrophotometric determination of quercetin with VO₂⁺. *Microchim. Acta.* 1979, 71(5-6), 365-370.
23. Alimarin I.P., Dorokhova E.N., Kazanskii I.V. i dr. Reaktsii geteropolikislot s osnovnymi krasitelyami i ikh primenenie v analize. *Zhurn. analit. khimii.* 1984, 34(6), 965-973.
24. Papaconstantinou E., Rore M.T. Heteropolyblues V. Electronic spectra one-to-six electron blues of 18-metaldiphosphate anions. *Inorg.Chem.* 1970, 9(3), 667-673.
25. Vishnikin A.B., Sklenarova H., Solich P., Petrushina G.A., Tsiganok L.P. Determination of ascorbic acid with Wells-Dawson type molybdophosphate in sequential injection system. *Anal. Lett.* 2011, 44(1-3), 514-527.
26. Kachan I.A., Zaporozhets O.A., Zin'ko L.S., Koval' A.A. Tverdogaznospektrofotometricheskoe opredelenie vosstanovitelei v rastvore po reaktsii obrazovaniya «sinej» geteropolikisloty. *Metody i ob"ekty khim. analiza.* 2006, 1(2), 127-131.
27. Sposib viznachennya zagal'nogo vmistu polifenoliv. Pat. № 101250. Ukraïna. MPK G01N 21/63, C07C 39/00, C07C 39/12. Zaporozhets' O.A., Bas Yu.P., Petrukh M.V. – № a 2011 10127; zayavleno 16.08.2011; nadr. 27.02.2012; Byul. № 4. – 6 s.
28. Alman M., Easton E.B. Selective determination of ascorbic acid with a novel hybrid material based 1-butyl-3-methylimidazolium tetrafluoroborate ionic liquid and the Dawson type ion [R₂Mo₁₈O₆₂]⁶⁻ immobilized on glassy carbon. *Electrochim. Acta.* 2011, 56(7), 2847-2855.
29. Lur'e Yu.Yu. *Spravochnik po analiticheskoi khimii.* Moskva: Khimiya, 1979, S. 278.
30. Timbola A.K., de Souza C.D., Giacomelli C., Spinelli A. Electrochemical oxidation of quercetin in hydro-alcoholic solution. *J. Braz. Chem. Soc.* 2006, 17(1), 139-148.
31. Tomson R.X. Struktura i reaktsionnaya sposobnost' fenol'nykh soedinenii. *Biokhimiya fenol'nykh soedinenii.* Pod red. Dzh. Kharborna. Moskva, 1968. S.233.
32. Ramešová Š., Sokolová R., Degano I., Bulíčková J., Žabka J., Gál M. On the stability of the bioactive flavonoids quercetin and luteolin under oxygen-free conditions. *Anal. Bioanal. Chem.* 2012, 402, 975-982.
33. Bulatov A.V., Petrova A.V., Vishnikin A.B., Moskvina A.L., Moskvina L.N. Stepwise injection spectrophotometric determination of epinephrine. *Talanta.* 2012, 96, 62-67.
34. Tsiganok L.P. Osobennosti reaktsii obrazovaniya i vosstanovleniya geteropolikompleksov elementov III-A gruppy Periodicheskoi sistemy.
35. L.P. Tsiganok, A.B. Vishnikin. Dnepropetrovsk: DGU, 1996. S.202.
36. Vishnikin A.B., Al-Shwaiyat M.K.E.A., Petrushina G.A., Tsiganok L.P., Andruch V., Bazel Ya.R., Sklenarova H., Solich P. Highly sensitive sequential injection determination of p-aminophenol in paracetamol formulations with
37. 18-molybdodiphosphate heteropoly anion based on elimination of Schlieren effect. *Talanta.* 2012, 96, 230-235.
38. Vishnikin A.B., Al-Shwaiyat M.K.E.A., Petrushina G.A., Tsiganok L.P., Andruch V., Bazel Ya.R., Sklenarova H., Solich P. Highly sensitive sequential injection determination of p-aminophenol in paracetamol formulations with
39. 18-molybdodiphosphate heteropoly anion based on elimination of Schlieren effect. *Talanta.* 2012, 96, 230-235.